

CDMO for PolyAmino Acids



Why PolyAmino Acids - PAA

- Functional Excipients
- Biodegradable Polymers
- Drug Delivery & Drug Conjugate
- Alternative to PEG (i.e. polysarcosine)
- Less Immunogenic Choice
- Half-life increase of proteins



A unique Integrated Solution

- Talented & Multidisciplinary R&D team
- Broad network for access cutting-edge technologies
- Integrated NCA supply (Phosgene expert)
- Ring Opening Polymerisation (ROP) technology
- Custom PAA synthesis & tailor-made
- From R&D to commercial supply



N°1 Producer of NCAs, 25+ years expertise



PMC Isochem services

- 45 years in large scale Manufacturing under GMP
- RSM, Intermediates, APIs and Excipients
- Analytical Development & QC
- Process development and scale up
- 3 FDA & EU qualified facilities (France)



Preclinical, clinical and commercial batches

- Lean approach to provide faster, better and cost effective solutions
- GMP batches for clinical trials
- Stability studies, process & analytical methods validations, reports...
- Full regulatory support (CMC-DMF Documentation)











PMC Isochem™ part of PMC Group - May 2022 - Michel Riehl - PMC Isochem - Design graphic : Kissagram Design

Why a PAA based delivery technology?

Non-exhaustive examples of PAA

linear, branched or grafted architectures

PAA HOMOPOLYMERS

Poly-L-amino acids

Poly-sarcosine


PAA COPOLYMERS

Methoxy-poly(ethylene glycol)-block-poly(L-amino acids)

Poly(L-amino acids)-block-poly(L-amino acids)

FUNCTIONALIZED PAA

Functionalized monoblock and diblock polyaminoacids



L or D amino acids configuration R, R' : amino acid side chains

R1 : alkyl group or functionalized alkyl group

R2: H or methyl group

R3: H or capping group

Growing interest for advanced & new polypeptides backbone

- PolyAmino Acids (PAAs) based materials have gained more interest in:
- > Drug delivery
- > Drug conjugates
- PAAs demonstrate remarkable biocompatibility and biodegradability due to the nature of the amino acid monomers building blocks
- · Less Immunogenic material and beneficial replacement of PEG
- · Use of activated amino acids monomers by ROP chemistry:
- > Narrow dirtibution of molar
- > Minimal side product formation
- > High reproducibility
- > Versatile architectures
- > Precise functionalisation of polypeptide backbone

O. Zagorodko et al, Macromol. Biosci. 2017

A global offer from amino acid to therapeutic solution

Amino acid

Amino acid-NCA Monomer

Polyamino acid

Drug Delivery / Biomaterials



Therapeutic areas

Oncology, Gene therapy, Immunotherapy, Ophtalmology, Regenerative medicine, Infectious and inflammatory diseases...



pmc 6 isochem

Your partner for Smart Delivery **Technologies**

GENERAL REQUESTS

Tel.: +33 1 64 99 0300

KARINE SALIM

k.salim@pmcisochem.fr

Tel.: +33 6 30 12 8461

